

Solutions to the Associated q -Askey-Wilson Polynomial Recurrence Relation *

Dharma P. Gupta and David R. Masson

Department of Mathematics
University of Toronto
Toronto Ontario M5S 1A1
Canada

In honour of Walter Gautschi on the occasion of his 65th birthday.

Abstract

A ${}_{10}\phi_9$ contiguous relation is used to derive contiguous relations for a very-well-poised ${}_8\phi_7$. These in turn yield solutions to the associated q -Askey-Wilson polynomial recurrence relation, expressions for the associated continued fraction, the weight function and a q -analogue of a generalized Dougall's theorem.

Key words and phrases: contiguous relations, Askey-Wilson polynomials, continued fraction, Pincherle's theorem, Jacobi matrix, spectrum, Dougall's theorem.

AMS subject classification: 33D45, 40A15, 39A10, 47B39.

* Research partially supported by NSERC (Canada).

1. Introduction.

A deeper insight into the properties of classical orthogonal polynomials is obtained by studying their associated cases. For recent work on associated Hermite, Laguerre, ultraspherical, Jacobi, continuous Hahn, continuous dual Hahn, big q -Laguerre and big q -Jacobi polynomials see [2], [4], [8], [9], [11], [12], [14], [15], [23]. The most general case of associated q -Askey-Wilson polynomials [16] and their $q \rightarrow 1$ limit, the associated Wilson polynomials, have also investigated [13], [18].

In this paper we examine the associated q -Askey-Wilson polynomial case using the methods of [18]. All of our parameters are, in general, complex. We essentially follow the notation in Gasper and Rahman [5], except that we omit the designation q for the base in the q -shifted factorials and basic hypergeometric functions. Thus we have, with $|q| < 1$,

$$(a)_0 := 1, \quad (a)_n := \prod_{j=1}^n (1 - aq^{j-1}), \quad n > 0, \text{ or } n = \infty,$$

$$(a_1, a_2, \dots, a_k)_n := \prod_{j=1}^k (a_j)_n.$$

The basic hypergeometric series ${}_r+1\phi_r$ is

$${}_r+1\phi_r \left(\begin{matrix} a_1, & a_2, & \dots, & a_{r+1} \\ b_1, & b_2, & \dots, & b_r \end{matrix} ; z \right) := \sum_{n=0}^{\infty} \frac{(a_1, a_2, \dots, a_{r+1})_n}{(b_1, b_2, \dots, b_r, q)_n} z^n.$$

The explicit form of monic q -Askey-Wilson polynomials is given by [1]:

$$(1.1) \quad P_n(z) = P_n(z; \alpha, \beta, \gamma, \delta) = \frac{(\alpha\beta, \alpha\gamma, \alpha\delta)_n (\alpha\beta\gamma\delta/q)_n}{(2\alpha)^n (\alpha\beta\gamma\delta/q)_{2n}} \\ \times {}_4\phi_3 \left(\begin{matrix} q^{-n}, \alpha\beta\gamma\delta q^{n-1}, \alpha u, \alpha/u \\ \alpha\beta, \alpha\gamma, \alpha\delta \end{matrix} ; q \right), \quad z = \frac{u + u^{-1}}{2}$$

which satisfy the three-term recurrence

$$(1.2) \quad P_{n+1}(z) - (z - a_n)P_n(z) + b_n^2 P_{n-1}(z) = 0, \\ a_n = a_n(\alpha, \beta, \gamma, \delta) = -A_n - B_n + \frac{\alpha}{2} + \frac{1}{2\alpha}, \\ b_n^2 = b_n^2(\alpha, \beta, \gamma, \delta) = A_{n-1}B_n,$$

$$A_n = \frac{(1 - \alpha\beta\gamma\delta q^{n-1})(1 - \alpha\beta q^n)(1 - \alpha\gamma q^n)(1 - \alpha\delta q^n)}{2\alpha(1 - \alpha\beta\gamma\delta q^{2n-1})(1 - \alpha\beta\gamma\delta q^{2n})},$$

$$B_n = \frac{\alpha(1 - q^n)(1 - \beta\gamma q^{n-1})(1 - \beta\delta q^{n-1})(1 - \gamma\delta q^{n-1})}{2(1 - \alpha\beta\gamma\delta q^{2n-2})(1 - \alpha\beta\gamma\delta q^{2n-1})}.$$

Associated monic q -Askey-Wilson polynomials are defined [16] by introducing an extra parameter, say ϵ , through a shift the discrete variable n in the coefficients of (1.2). Here we choose to replace n by $n + (\log \epsilon)/(\log q)$ so that q^n is replaced by ϵq^n . The new polynomial set $\{P_n(z; \epsilon)\}$ is an initial value solution ($X_{-1} = P_{-1} = 0$, $X_0 = P_0 = 1$) to the difference equation

$$(1.3) \quad X_{n+1} - (z - a_v)X_n + b_v^2 X_{n-1} = 0 \quad v := n + (\log \epsilon)/(\log q).$$

The associated q -Askey-Wilson polynomials have been studied by Ismail and Rahman [16]. They derive contiguous relations for a very well poised ${}_8\phi_7$ series and then obtain two linearly independent solutions of the three-term recurrence relation for the associated Askey-Wilson polynomials. Explicit representations of two families of associated Askey-Wilson polynomials are given. Corresponding continued fractions are identified and weight functions obtained.

On the other hand Masson [18] studied the associated Wilson polynomials by obtaining ${}_7F_6$ contiguous relations with a procedure (see Wilson [22]) different from that employed by Ismail and Rahman. He discussed explicit forms of solutions to the three-term recurrence satisfied by associated Wilson polynomials and identified minimal solutions for different regions. Using Pincherle's theorem [6],[17], he obtained a number of new continued fraction representations and discussed the spectral properties of the corresponding Jacobi matrix. A weight function was obtained and his method gave a generalization of Dougall's theorem [3] which removed both the terminating and the balancing conditions.

The object of the present paper is to study the associated q -Askey-Wilson polynomials using Masson's approach [18]. In spite of the fact that some of the basic results here will overlap with the results obtained by Ismail and Rahman [16], we have to offer a different procedure for obtaining the ${}_8\phi_7$ contiguous relations, the minimal solution to the associated Askey-Wilson polynomial recurrence, the corresponding continued fraction

representations and an extension of Masson's generalization of Dougall's theorem [18] to the basic hypergeometric case. Moreover we again see, as in many previous cases ([7], [8], [9], [18], [19], [21]), that Pincherle's theorem [6],[17] plays a crucial role. The importance of Pincherle's theorem was re-emphasized by Gautschi in 1967 [6].

In Section 2, following Wilson's method [22], we derive a pair of three-term recurrence relations for very-well-poised balanced and terminating $_{10}\phi_9$ basic hypergeometric functions. From these recurrence relations and using certain known transformations [5] we obtain a number of explicit solutions to (1.3). Minimal solutions for different regions of the complex plane are identified. In Section 3, the associated continued fractions are obtained. In Section 4 we derive an expression for the weight function of orthogonality and a Wronskian identity which turns out to be the basic analogue of a generalized Dougall's theorem [18].

2. Recurrence relation and solutions.

In a recent paper [10] we have derived contiguous relations for a very- well-poised balanced $_{10}\phi_9$ basic hypergeometric function

$$(2.1) \quad \phi := {}_{10}\phi_9 \left(\begin{matrix} a, q\sqrt{a}, -q\sqrt{a}, b, c, d, e, f, g, h \\ \sqrt{a}, -\sqrt{a}, \frac{aq}{b}, \frac{aq}{c}, \dots, \frac{aq}{h} \end{matrix} ; q \right)$$

where one of the numerator parameters, say h , is equal to q^{-n} , $n = 0, 1, \dots$, $bcdefgh = a^3q^2$ and $|q| < 1$. One such relation is

$$(2.2) \quad \begin{aligned} & \frac{b(1-c)(1-\frac{a}{c})(1-\frac{aq}{c})(1-\frac{aq}{bd})(1-\frac{aq}{be})(1-\frac{aq}{bf})(1-\frac{aq}{bg})(1-\frac{aq}{bh})}{(1-\frac{cq}{b})} \\ & \quad \times [\phi(b-, c+) - \phi] \\ & - \frac{c(1-b)(1-\frac{a}{b})(1-\frac{aq}{b})(1-\frac{aq}{cd})(1-\frac{aq}{ce})(1-\frac{aq}{cf})(1-\frac{aq}{cg})(1-\frac{aq}{ch})}{(1-\frac{bq}{c})} \\ & \quad \times [\phi(c-, b+) - \phi] \\ & - \frac{aq}{c}(1-\frac{c}{b})(1-\frac{bc}{aq})(1-d)(1-e)(1-f)(1-g)(1-h)\phi = 0. \end{aligned}$$

Note that we are using the notation $\phi(b-, c+)$ to mean ϕ in (2.1) with b replaced by b/q and c replaced by cq .

In (2.2), if we write $h = q^{-n}$, $g = a^3 q^{n+2}/bcdef$ and let $n \rightarrow \infty$, we obtain

$$(2.3) \quad \frac{bcdef}{a^2 q} \frac{(1-c)(1-\frac{a}{c})(1-\frac{aq}{c})(1-\frac{aq}{bd})(1-\frac{aq}{be})(1-\frac{aq}{bf})}{(1-\frac{cq}{b})} [{}_8\phi_7(b-, c+) - {}_8\phi_7] \\ - \frac{bcdef}{a^2 q} \frac{(1-b)(1-\frac{a}{b})(1-\frac{aq}{b})(1-\frac{aq}{cd})(1-\frac{aq}{ce})(1-\frac{aq}{cf})}{(1-\frac{bq}{c})} [{}_8\phi_7(c-, b+) - {}_8\phi_7] \\ - \frac{aq}{c} (1-\frac{c}{b})(1-\frac{bc}{aq})(1-d)(1-e)(1-f) {}_8\phi_7 = 0,$$

where

$${}_8\phi_7 = {}_8\phi_7 \left(\begin{matrix} a, q\sqrt{a}, -q\sqrt{a}, b, c, d, e, f \\ \sqrt{a}, -\sqrt{a}, \frac{aq}{b}, \frac{aq}{c}, \frac{aq}{d}, \frac{aq}{e}, \frac{aq}{f} \end{matrix} ; \frac{a^2 q^2}{bcdef} \right) \\ = W(a; b, c, d, e, f) \quad \text{say}.$$

For non-terminating ${}_8\phi_7$, the convergence condition is $|a^2 q^2 / bcdef| < 1$. In (2.3) we now put

$$a = \frac{\alpha\beta\gamma u}{q}, \quad b = q^{-n}/\epsilon, \quad c = \epsilon s q^{n-1}, \quad d = \alpha u, \quad e = \beta u, \quad f = \gamma u, \\ s = \alpha\beta\gamma\delta.$$

Renormalizing and simplifying, we obtain the associated Askey-Wilson recurrence

$$(2.5) \quad X_{n+1}^{(1)} - (z - a'_n)X_n^{(1)} + b_n'^2 X_{n-1}^{(1)} = 0$$

with

$$z = \frac{u + u^{-1}}{2}, \\ a'_n = -A'_n - B'_n + \frac{\alpha}{2} + \frac{1}{2\alpha} \\ b_n'^2 = A'_{n-1} B'_n \\ A'_n = \frac{(1 - sq^{n-1}\epsilon)(1 - \alpha\beta\epsilon q^n)(1 - \alpha\gamma\epsilon q^n)(1 - \alpha\delta\epsilon q^n)}{2\alpha(1 - sq^{2n-1}\epsilon^2)(1 - sq^{2n}\epsilon^2)}, \\ B'_n = \frac{\alpha(1 - \epsilon q^n)(1 - \beta\gamma\epsilon q^{n-1})(1 - \beta\delta\epsilon q^{n-1})(1 - \gamma\delta\epsilon q^{n-1})}{2(1 - s\epsilon^2 q^{2n-2})(1 - s\epsilon^2 q^{2n-1})}$$

and a solution of (2.5) is therefore

$$(2.6) \quad X_n^{(1)}(u) = \left(\frac{u}{2}\right)^n \frac{(sueq^n/\delta)_\infty (s\epsilon^2 q^{2n-1})_\infty}{(seq^{n-1})_\infty (\delta eq^n/u)_\infty (\alpha\beta eq^n, \alpha\gamma eq^n, \beta\gamma eq^n)_\infty} \\ \times W\left(\frac{\alpha\beta\gamma u}{q}; \frac{q^{-n}}{\epsilon}, \epsilon s q^{n-1}, \alpha u, \beta u, \gamma u\right),$$

with convergence condition $|q/\delta u| < 1$.

The solution $X_n^{(1)}$ is a generalization of the Askey-Wilson solution (1.1) to the $\epsilon = 1$ case of (2.5). This can be verified through an application of Watson's formula ([5], III.18, p.242) connecting a terminating ${}_4\phi_3$ with an ${}_8\phi_7$.

A second linearly independent solution of (2.5) may be obtained with the help of the reflection symmetry transformation $v \rightarrow -v - 1$, $(\alpha, \beta, \gamma, \delta) \rightarrow (q/\alpha, q/\beta, q/\gamma, q/\delta)$. Using (see also [10])

$$b_{-v-1}^2(q/\alpha, q/\beta, q/\gamma, q/\delta) = b_{v+1}^2(\alpha, \beta, \gamma, \delta) \\ a_{-v-1}(q/\alpha, q/\beta, q/\gamma, q/\delta) = a_v(\alpha, \beta, \gamma, \delta)$$

and renormalizing, we arrive at the solution

$$(2.7) \quad X_n^{(2)}(u) = \left(\frac{u}{2}\right)^n \frac{(s\epsilon^2 q^{2n-1})_\infty (\epsilon\delta u q^{n+1})_\infty}{(\epsilon q^{n+1})_\infty (\beta\delta eq^n, \gamma\delta eq^n, \alpha\delta eq^n)_\infty (\frac{s\epsilon}{\delta u} q^{n-1})_\infty} \\ \times W\left(\frac{q^2 u}{\alpha\beta\gamma}; \epsilon q^{n+1}, \frac{q^{-n+2}}{\epsilon s}, \frac{qu}{\alpha}, \frac{qu}{\beta}, \frac{qu}{\gamma}\right),$$

with convergence condition $|\delta/u| < 1$.

In order to obtain a third solution of (2.5) we make the following alterations in (2.2): Interchange c and h , replace $a, b, c, \dots, g$ by $h^2/a, hb/a, hc/a, \dots, hg/a$ respectively, put $h = q^{-n}$, n being a positive integer, and reverse the ${}_{10}\phi_9$ series. This gives

$$(2.8) \quad \frac{\frac{b}{a}(1-h)(1-\frac{h}{a})(1-\frac{hq}{a})(1-\frac{aq}{bd})(1-\frac{aq}{be})(1-\frac{aq}{bf})(1-\frac{aq}{bg})(1-\frac{aq}{bc})}{(1-\frac{aq}{b})} \\ \times [c_2\phi_+(b-) - c_1\phi] \\ - \frac{(1-\frac{bh}{a})(1-\frac{h}{b})(1-\frac{hq}{b})(1-\frac{q}{d})(1-\frac{q}{e})(1-\frac{q}{f})(1-\frac{q}{g})(1-\frac{q}{c})}{(1-\frac{bq}{a})} \\ \times [c_3\phi_-(b+) - c_1\phi] \\ - \frac{q}{a}(1-\frac{a}{b})(1-\frac{b}{q})(1-\frac{hd}{a})(1-\frac{he}{a})(1-\frac{hf}{a})(1-\frac{hg}{a})(1-\frac{hc}{a})c_1\phi = 0,$$

where

$$\begin{aligned}
c_1 &= \frac{(\sqrt{a})_n (-\sqrt{a})_n \left(\frac{aq}{b}\right)_n \left(\frac{aq}{c}\right)_n \cdots \left(\frac{aq}{g}\right)_n (aq^{n+1})_n}{(a)_n (q\sqrt{a})_n (-q\sqrt{a})_n (b)_n (c)_n \cdots (g)_n} (-1)^n q^{n(n-1)/2} \\
c_2 &= \frac{(q\sqrt{a})_{n-1} (-q\sqrt{a})_{n-1} \left(\frac{aq^3}{b}\right)_{n-1} \left(\frac{aq^2}{c}\right)_{n-1} \cdots \left(\frac{aq^2}{g}\right)_{n-1} (aq^{n+2})_{n-1}}{(aq^2)_{n-1} (q^2\sqrt{a})_{n-1} (-q^2\sqrt{a})_{n-1} (b)_{n-1} (cq)_{n-1} \cdots (gq)_{n-1}} (-1)^{n-1} q^{(n-1)(n-2)/2} \\
c_3 &= \frac{\left(\frac{\sqrt{a}}{q}\right)_{n+1} \left(-\frac{\sqrt{a}}{q}\right)_{n+1} \left(\frac{a}{bq}\right)_{n+1} \left(\frac{a}{c}\right)_{n+1} \cdots \left(\frac{a}{g}\right)_{n+1} (aq^n)_{n+1}}{\left(\frac{a}{q^2}\right)_{n+1} (\sqrt{a})_{n+1} (-\sqrt{a})_{n+1} (b)_{n+1} \left(\frac{c}{q}\right)_{n+1} \cdots \left(\frac{g}{q}\right)_{n+1}} (-1)^{n+1} q^{n(n+1)/2}.
\end{aligned}$$

Note that $\phi_+(b-)$ and $\phi_-(b+)$ refer to ϕ with (a, c, d, e, f, g, h) replaced by $(aq^2, cq, dq, eq, fq, gq, hq)$ and $(aq^{-2}, c/q, d/q, e/q, f/q, g/q, h/q)$ respectively.

Taking the limit $h = q^{-n} \rightarrow \infty$, $g = a^3 q^{2+n}/bcdef \rightarrow 0$, we obtain

$$\begin{aligned}
(2.9) \quad & \frac{b(1 - \frac{aq}{bd})(1 - \frac{aq}{be})(1 - \frac{aq}{bf})(1 - \frac{aq}{bc})}{a(1 - \frac{aq}{b})} \\
& \times \left[\frac{a^2 q^2}{bcdef} \frac{(1 - aq)(1 - aq^2)(1 - c)(1 - d)(1 - e)(1 - f)}{(1 - \frac{aq}{b})(1 - \frac{aq^2}{b})(1 - \frac{aq}{c})(1 - \frac{aq}{d})(1 - \frac{aq}{e})(1 - \frac{aq}{f})} W_+(b-) - W \right] \\
& - \frac{(1 - \frac{q}{d})(1 - \frac{q}{e})(1 - \frac{q}{f})(1 - \frac{q}{c})}{(1 - \frac{bq}{a})} \\
& \times \left[\frac{bcdef}{a^2 q^2} \frac{(1 - \frac{a}{bq})(1 - \frac{a}{b})(1 - \frac{a}{c})(1 - \frac{a}{d})(1 - \frac{a}{e})(1 - \frac{a}{f})}{(1 - \frac{a}{q})(1 - a)(1 - \frac{c}{q})(1 - \frac{d}{q})(1 - \frac{e}{q})(1 - \frac{f}{q})} W_-(b+) - W \right] \\
& - \frac{q}{a} \left(1 - \frac{a}{b}\right) \left(1 - \frac{b}{q}\right) \left(1 - \frac{a^2 q^2}{bcdef}\right) W = 0.
\end{aligned}$$

In (2.9), writing

$$\begin{aligned}
(2.10) \quad & a = \frac{\beta\gamma\delta\epsilon^2}{u} q^{2n}, \quad b = \frac{q}{\alpha u}, \quad c = q^{n+1}\epsilon, \quad d = \beta\delta\epsilon q^n \\
& e = \gamma\delta\epsilon q^n, \quad f = \beta\gamma\epsilon q^n, \quad s = \alpha\beta\gamma\delta
\end{aligned}$$

and renormalizing, we obtain a third solution to (2.5) given by

$$\begin{aligned}
(2.11) \quad & X_n^{(3)}(u) = \frac{1}{(2u)^n} \\
& \times \frac{(s\epsilon^2 q^{2n})_\infty (s\epsilon^2 q^{2n-1})_\infty \left(\frac{\beta\gamma\delta\epsilon}{u} q^n\right)_\infty \left(\frac{\epsilon\beta q^{n+1}}{u}, \frac{\epsilon\gamma q^{n+1}}{u}, \frac{\epsilon\delta q^{n+1}}{u}\right)_\infty}{(\epsilon q^{n+1})_\infty (\epsilon s q^{n-1})_\infty (\alpha\beta\epsilon q^n, \alpha\gamma\epsilon q^n, \alpha\delta\epsilon q^n, \beta\gamma\epsilon q^n, \beta\delta\epsilon q^n, \gamma\delta\epsilon q^n)_\infty \left(\frac{\beta\gamma\delta\epsilon^2}{u} q^{2n+1}\right)_\infty} \\
& \times W\left(\frac{\beta\gamma\delta\epsilon^2}{u} q^{2n}; \frac{q}{\alpha u}, \epsilon q^{n+1}, \beta\delta\epsilon q^n, \gamma\delta\epsilon q^n, \beta\gamma\epsilon q^n\right),
\end{aligned}$$

with convergence condition $|\alpha/u| < 1$. A transformation ([5], III.23, p.243) may be applied to $X_n^{(3)}$ to obtain this solution in a form which is symmetric in the parameters $\alpha, \beta, \gamma, \delta$. We get the solution (omitting constant factors)

$$(2.12) \quad X_n^{(4)}(u) = \frac{1}{(2u)^n} \frac{(\epsilon^2 s q^{2n-1})_\infty \left(\frac{\alpha\epsilon}{u} q^{n+1}, \frac{\beta\epsilon}{u} q^{n+1}, \frac{\gamma\epsilon}{u} q^{n+1}, \frac{\delta\epsilon}{u} q^{n+1}\right)_\infty}{(\epsilon q^{n+1})_\infty \left(\frac{q^{n+2}\epsilon}{u^2}\right)_\infty (\alpha\beta\epsilon q^n, \alpha\delta\epsilon q^n, \alpha\gamma\epsilon q^n, \beta\gamma\epsilon q^n, \beta\delta\epsilon q^n, \gamma\delta\epsilon q^n)_\infty} \\ \times W\left(\frac{q^{n+1}\epsilon}{u^2}; \epsilon q^{n+1}, \frac{q}{\alpha u}, \frac{q}{\beta u}, \frac{q}{\gamma u}, \frac{q}{\delta u}\right)$$

where the convergence condition is $|s q^{n-1}\epsilon| < 1$.

The large n behaviour of $X_n^{(4)}(u)$ is easily seen to be

$$X_n^{(4)}(u) \stackrel{n \rightarrow \infty}{\sim} \frac{1}{(2u)^n}$$

while the corresponding solution

$$X_n^{(4)}(1/u) \sim \left(\frac{u}{2}\right)^n.$$

It is evident that $X_n^{(4)}(u)$ is a sub-dominant or minimal solution for $|u| > 1$, $|s q^{n-1}\epsilon| < 1$ while $X_n^{(4)}(1/u)$ is sub-dominant (minimal) for $|u| < 1$, $|s q^{n-1}\epsilon| < 1$.

We proceed to get another solution by making the following parameter replacements in (2.9):

$$(2.14) \quad \begin{aligned} a &= \frac{q^{-2n+1}}{\beta\gamma\delta u\epsilon^2}, & b &= \frac{\alpha}{u}, & c &= \frac{q^{-n}}{\epsilon}, & d &= \frac{1}{\beta\delta\epsilon} q^{-n+1}, \\ e &= \frac{1}{\gamma\delta\epsilon} q^{-n+1}, & f &= \frac{1}{\beta\gamma\epsilon} q^{-n+1}. \end{aligned}$$

Note that this is just the reflection transformation that was used to obtain the solution $X_n^{(2)}$ from $X_n^{(1)}$ but now it is being applied to $X_n^{(3)}$.

After renormalization, omitting constant factors and using the transformation [5, III.23, p.243] we arrive at the following solution

$$(2.15) \quad X_n^{(5)}(u) = \frac{1}{(2u)^n} \frac{(s q^{2n-1}\epsilon^2)_\infty (u^2\epsilon q^n)_\infty}{(s q^{n-1}\epsilon)_\infty (\alpha u\epsilon q^n, \beta u\epsilon q^n, \gamma u\epsilon q^n, \delta u\epsilon q^n)_\infty} \\ \times W\left(\frac{q^{-n}}{\epsilon u^2}; \frac{q^{-n}}{\epsilon}, \frac{\alpha}{u}, \frac{\beta}{u}, \frac{\gamma}{u}, \frac{\delta}{u}\right)$$

where the convergence condition is $|q^{-n+2}/s\epsilon| < 1$.

Note that, when $\epsilon = 1$, the solution (2.15) is proportional to the Askey-Wilson polynomial $P_n(z; \alpha, \beta, \gamma, \delta)$ in (1.1) and can be used to give $P_n(z)$ in a form which is explicitly symmetric in its parameters. That is

$$P_n(z) = (2u)^{-n} \frac{(\alpha u, \beta u, \gamma u, \delta u, s/q)_n}{(s/q)_{2n} (u^2)_n} W(q^{-n}/u^2; q^{-n}, \frac{\alpha}{u}, \frac{\beta}{u}, \frac{\gamma}{u}, \frac{\delta}{u}).$$

A solution in which the convergence condition is independent of z and also independent of the parameters $\alpha, \beta, \gamma, \delta$ can be obtained from (2.12) by applying the transformation ([5], III.24, p.243). This gives

$$(2.16) \quad X_n^{(6)}(u) = \frac{1}{(2u)^n} \frac{(s\epsilon^2 q^{2n-1})_\infty (s\epsilon q^n/\alpha u, s\epsilon q^n/\beta u, s\epsilon q^n/\gamma u, s\epsilon q^n/\delta u)_\infty}{(\epsilon s q^{n-1})_\infty (s\epsilon q^n/u^2)_\infty (\alpha\beta\epsilon q^n, \alpha\gamma\epsilon q^n, \alpha\delta\epsilon q^n, \beta\gamma\epsilon q^n, \beta\delta\epsilon q^n, \gamma\delta\epsilon q^n)_\infty} \\ \times W\left(\frac{s\epsilon q^{n-1}}{u^2}; \epsilon s q^{n-1}, \frac{\alpha}{u}, \frac{\beta}{u}, \frac{\gamma}{u}, \frac{\delta}{u}\right)$$

with convergence condition $|q^{n+1}\epsilon| < 1$.

Since $X_n^{(6)}(u) \stackrel{n \rightarrow \infty}{\sim} \frac{1}{(2u)^n}$, we have that $X_n^{(6)}(u)$ is a sub-dominant(minimal) solution for $|u| > 1$ while $X_n^{(6)}(1/u)$ is a sub-dominant(minimal) solution for $|u| < 1$. Note that $X_n^{(6)}$ is really the same solution as $X_n^{(4)}$, but analytically continued.

Summarizing we have

Theorem 1. *The associated Askey-Wilson equation (2.5) has solutions $X_n^{(i)}(u)$ and $X_n^{(i)}(1/u)$, $i = 1, 2, \dots, 6$ given by (2.6), (2.7), (2.11), (2.12), (2.15) and (2.16). If $|u| > 1$, then $X_n^{(3)}(u)$, $X_n^{(4)}(u)$ and $X_n^{(6)}(u)$ (which are connected by general ${}_8\phi_7$ transformations) each represent a minimal solution $X_n^{(s)}(z)$ for the parameter values $|\frac{\alpha}{u}| < 1$, $|\epsilon s q^{n-1}| < 1$, $|\epsilon q^{n+1}| < 1$ respectively.*

3. Continued fraction representations.

The continued fraction associated with the difference equation (2.5) is

$$(3.1) \quad CF(z) := z - a'_0 + \cfrac{\infty}{K} \left(\cfrac{-b_n'^2}{z - a'_n} \right).$$

If $b_n'^2 \neq 0$, $n \geq 1$, by Pincherle's theorem [6], [17] and Theorem 1 of Section 2 we have

$$(3.2) \quad \cfrac{1}{CF(z)} = \cfrac{X_0^{(s)}(z)}{b_0'^2 X_{-1}^{(s)}(z)}.$$

Using $X_0^{(s)}(z) = X_0^{(4)}(u)$, $X_{-1}^{(s)}(z) = X_{-1}^{(4)}(u)$, $|u| > 1$, $z = \frac{u+u^{-1}}{2}$, we have from (2.12),

$$(3.3) \quad \begin{aligned} \cfrac{1}{CF(z)} &= \cfrac{2}{u} \cfrac{(1 - \epsilon q/u^2)(1 - s\epsilon^2/q^2)(1 - s\epsilon^2/q)}{(1 - \alpha\epsilon/u)(1 - \beta\epsilon/u)(1 - \gamma\epsilon/u)(1 - \delta\epsilon/u)(1 - s\epsilon/q^2)} \\ &\times \cfrac{W\left(q\epsilon/u^2; q\epsilon, q/\alpha u, q/\beta u, q/\gamma u, q/\delta u\right)}{W\left(\epsilon/u^2; \epsilon, q/\alpha u, q/\beta u, q/\gamma u, q/\delta u\right)} \end{aligned}$$

for $|u| > 1$ and $|s\epsilon/q^2| < 1$. For $|u| < 1$, $|s\epsilon/q^2| < 1$ we replace u in (3.3) by $1/u$.

Other representations for the continued fraction for different parameter ranges are obtained by taking a different representation for the minimal solution. For example, by taking $X_0^{(s)}(z) = X_0^{(6)}(u)$ and $X_{-1}^{(s)}(z) = X_{-1}^{(6)}(u)$, $|u| > 1$, we obtain the continued fraction representation

$$(3.4) \quad \begin{aligned} \cfrac{1}{CF(z)} &= \cfrac{2}{u} \cfrac{(1 - s\epsilon^2/q^2)(1 - s\epsilon^2/q)(1 - s\epsilon/u^2 q)}{(1 - \epsilon)(1 - s\epsilon/\alpha u q)(1 - s\epsilon/\beta u q)(1 - s\epsilon/\gamma u q)(1 - s\epsilon/\delta u q)} \\ &\times \cfrac{W\left(\epsilon s/u^2 q; \epsilon s/q, \alpha/u, \beta/u, \gamma/u, \delta/u\right)}{W\left(s\epsilon/u^2 q^2; \epsilon s/q^2, \alpha/u, \beta/u, \gamma/u, \delta/u\right)}, \end{aligned}$$

for $|u| > 1$, $|\epsilon| < 1$.

4. Weight function.

We have seen above that a minimal solution exists for $|u| > 1$ and $|u| < 1$. There is no minimal solution for $z \in [-1, 1]$, ($|u| = 1$), where we have a continuous spectrum for the associated tridiagonal Jacobi matrix J with diagonal $(a'_0, a'_1, \dots)$ and $(b'_1, b'_2, \dots)$ above and below the diagonal. For a probability measure $d\omega(x; \epsilon)$ we have, in the case of real orthogonality ($b'_{n+1} > 0, a'_n$ real, $n \geq 0$) for the associated monic q -Askey-Wilson polynomials $P_n(x; \epsilon)$,

$$(4.1) \quad \int_{\mathbb{R}} P_n(x; \epsilon) P_m(x; \epsilon) d\omega(x; \epsilon) = \delta_{nm} \prod_{k=1}^n b'_k{}^2.$$

From [21], we have the representation

$$(4.2) \quad \frac{1}{CF(z)} = \int_{\mathbb{R}} \frac{d\omega(x; \epsilon)}{z - x} = \frac{X_0^{(s)}(z)}{b_0'^2 X_{-1}^{(s)}(z)}$$

and for the absolutely continuous part for $x \in [-1, 1]$,

$$(4.3) \quad \begin{aligned} \frac{d\omega(x; \epsilon)}{dx} &= \frac{1}{2\pi i b_0'^2} \left(\frac{X_0^{(s)}(x - i0)}{X_{-1}^{(s)}(x - i0)} - \frac{X_0^{(s)}(x + i0)}{X_{-1}^{(s)}(x + i0)} \right) \\ &= \frac{1}{2\pi i b_0'^2} \frac{\mathcal{W}(X_{-1}^{(s)}(x + i0), X_{-1}^{(s)}(x - i0))}{|X_{-1}^{(s)}(x - i0)|^2}, \end{aligned}$$

where

$$W(X_n, Y_n) := X_n Y_{n+1} - X_{n+1} Y_n.$$

From (2.5) we have

$$(4.4) \quad \mathcal{W}(X_{-1}^{(4)}(u), X_{-1}^{(4)}(1/u)) = \lim_{n \rightarrow \infty} \frac{\mathcal{W}(X_n^{(4)}(u), X_n^{(4)}(1/u))}{\prod_{k=0}^n b'_k{}^2}.$$

Using (2.5) and (2.12) we then obtain

$$(4.5) \quad \begin{aligned} &\mathcal{W}(X_{-1}^{(4)}(u), X_{-1}^{(4)}(1/u)) \\ &= 2(u - u^{-1}) \frac{\left(\frac{s\epsilon^2}{q^3}\right)_{\infty} \left(\frac{s\epsilon^2}{q^2}\right)_{\infty}}{\left(\frac{\alpha\beta\epsilon}{q}, \frac{\alpha\gamma\epsilon}{q}, \frac{\alpha\delta\epsilon}{q}, \frac{\beta\gamma\epsilon}{q}, \frac{\beta\delta\epsilon}{q}, \frac{\gamma\delta\epsilon}{q}, \frac{s\epsilon}{q^2}, \epsilon\right)_{\infty}}. \end{aligned}$$

Thus (4.3) with

$$X_n^{(s)}(x \pm i0) = X_n^{(4)}(e^{\mp i\theta}), \quad x = \cos \theta, \quad u = e^{i\theta}$$

gives for $-1 \leq x \leq 1$, $|s\epsilon/q^2| < 1$,

$$(4.6) \quad \frac{d\omega(x; \epsilon)}{dx} = \frac{2\sqrt{1-x^2}}{\pi} \frac{(1-s\epsilon^2/q)(1-s\epsilon^2/q^2)^2}{(1-s\epsilon/q^2)} \\ \times \frac{(\epsilon q/u^2, \epsilon q u^2, \alpha\beta\epsilon, \alpha\gamma\epsilon, \alpha\delta\epsilon, \beta\gamma\epsilon, \beta\delta\epsilon, \gamma\delta\epsilon, \epsilon q)_\infty}{(\alpha\epsilon/u, \alpha\epsilon u, \beta\epsilon/u, \beta\epsilon u, \gamma\epsilon/u, \gamma\epsilon u, \delta\epsilon/u, \delta\epsilon u, s\epsilon/q^2)_\infty} \\ \times \frac{1}{\left| W(\epsilon/u^2; q/\alpha u, q/\beta u, q/\gamma u, q/\delta u, \epsilon) \right|^2}.$$

This checks with the weight function obtained by Ismail and Rahman ([16], (4.31), p.218).

When $\epsilon = 1$, this reduces to the Askey-Wilson weight function [1]. Also from the positivity of the denominators in (3.4) there is no discrete spectrum if $-1 < \epsilon < 1$, $|\alpha|, |\beta|, |\gamma|, |\delta| < |q|^{1/2}$, [16]. Other conditions for the absence of a discrete spectrum may be deduced from (3.3) and (3.4). For example, from (3.3), there is no discrete spectrum if $-|q^2/s| < \epsilon < |q^2/s|$, $|\alpha|, |\beta|, |\gamma|, |\delta| > |q|^{1/2}$.

Next we use (4.3) with

$$\frac{d\omega(x; \epsilon)}{dx} = \frac{1}{2\pi i b_0^2} \left(\frac{X_0^{(6)}(u)}{X_{-1}^{(4)}(u)} - \frac{X_0^{(6)}(1/u)}{X_{-1}^{(4)}(1/u)} \right), \quad x = \cos \theta, \quad u = e^{i\theta}.$$

The right side simplifies to

$$(4.7) \quad \frac{1}{2\pi i} \frac{(1-s\epsilon^2/q^2)(1-s\epsilon^2/q)(\epsilon q)_\infty}{(s\epsilon/q^2)_\infty} \\ \times \left[2u \frac{(\epsilon q u^2)_\infty \left(\frac{s\epsilon}{\alpha} u, \frac{s\epsilon}{\beta} u, \frac{s\epsilon}{\gamma} u, \frac{s\epsilon}{\delta} u \right)_\infty W\left(\frac{s u^2 \epsilon}{q}; \frac{s\epsilon}{q}, \alpha u, \beta u, \gamma u, \delta u\right)}{(s u^2 \epsilon)_\infty (\alpha u \epsilon, \beta u \epsilon, \gamma u \epsilon, \delta u \epsilon)_\infty W\left(\epsilon u^2; \epsilon, \frac{q u}{\alpha}, \frac{q u}{\beta}, \frac{q u}{\gamma}, \frac{q u}{\delta}\right)} \right. \\ \left. - \frac{2(\epsilon q/u^2)_\infty (s\epsilon/\alpha u, s\epsilon/\beta u, s\epsilon/\gamma u, s\epsilon/\delta u)_\infty W\left(\frac{s}{u^2} \frac{\epsilon}{q}; \frac{s\epsilon}{q}, \frac{\alpha}{u}, \frac{\beta}{u}, \frac{\gamma}{u}, \frac{\delta}{u}\right)}{\left(\frac{s\epsilon}{u^2}\right)_\infty \left(\frac{\alpha\epsilon}{u}, \frac{\beta\epsilon}{u}, \frac{\gamma\epsilon}{u}, \frac{\delta\epsilon}{u}\right)_\infty W\left(\frac{\epsilon}{u^2}; \epsilon, \frac{q}{\alpha u}, \frac{q}{\beta u}, \frac{q}{\gamma u}, \frac{q}{\delta u}\right)} \right].$$

Equating (4.6) and (4.7), and writing

$$(4.8) \quad G(\alpha, \beta, \gamma, \delta, \epsilon, u) := \frac{1}{u} \frac{\left(\frac{\epsilon q}{u^2}\right)_\infty \left(\frac{s\epsilon}{\alpha u}, \frac{s\epsilon}{\beta u}, \frac{s\epsilon}{\gamma u}, \frac{s\epsilon}{\delta u}\right)_\infty}{\left(s\epsilon/u^2\right)_\infty \left(\frac{\alpha\epsilon}{u}, \frac{\beta\epsilon}{u}, \frac{\gamma\epsilon}{u}, \frac{\delta\epsilon}{u}\right)_\infty} \\ \times W\left(\frac{s\epsilon}{u^2 q}; \frac{s\epsilon}{q}, \frac{\alpha}{u}, \frac{\beta}{u}, \frac{\gamma}{u}, \frac{\delta}{u}\right) W\left(\epsilon u^2; \epsilon, \frac{q u}{\alpha}, \frac{q u}{\beta}, \frac{q u}{\gamma}, \frac{q u}{\delta}\right),$$

we obtain the identity

$$(4.9) \quad G(\alpha, \beta, \gamma, \delta, \epsilon, u) - G(\alpha, \beta, \gamma, \delta, \epsilon, 1/u) = (1/u - u) \frac{(1 - s\epsilon^2/q^2)}{(1 - s\epsilon/q^2)} \\ \times \frac{(\alpha\beta\epsilon, \alpha\gamma\epsilon, \alpha\delta\epsilon, \beta\gamma\epsilon, \beta\delta\epsilon, \gamma\delta\epsilon, \epsilon q/u^2, \epsilon q u^2)_\infty}{(\alpha\epsilon/u, \alpha\epsilon u, \beta\epsilon/u, \beta\epsilon u, \gamma\epsilon/u, \gamma\epsilon u, \delta\epsilon/u, \delta\epsilon u)_\infty}$$

which is a q -analogue of Masson's generalization of Dougall's theorem [18]. $\epsilon = 1$ gives a q -analogue of Dougall's theorem (see [3]).

We can recover from (4.9) the following identity which we had obtained in our earlier paper ([7], (31), p.723) for $\epsilon = 1$, $s = q^m$; $m = 1, 2, \dots$, i.e.

$$(4.10) \quad \left(\frac{q}{\alpha}\right)^{m-3} \left(\frac{1}{u} - u\right) \left[u^{m-2} \Pi_1(u) \Pi_2\left(\frac{1}{u}\right) - u^{2-m} \Pi_1\left(\frac{1}{u}\right) \Pi_2(u) \right] \\ = (\alpha\beta q^{-1}, \alpha\gamma q^{-1}, \alpha\delta q^{-1})_\infty \left(\frac{q^2}{\alpha\beta}, \frac{q^2}{\alpha\gamma}, \frac{q^2}{\alpha\delta}\right)_\infty \left(u^2, \frac{1}{u^2}\right)_\infty$$

for $m = 1, 2, \dots$ and $|u| = 1$, where

$$\Pi_1(u) = \left(\frac{q}{\alpha u}, \frac{q}{\beta u}, \frac{q}{\gamma u}, \frac{q}{\delta u}\right)_\infty \\ \Pi_2(u) = \left(\frac{\alpha}{u}, \frac{\beta}{u}, \frac{\gamma}{u}, \frac{\delta}{u}\right)_\infty.$$

References

1. R. Askey and J. Wilson, Some basic hypergeometric orthogonal polynomials that generalize Jacobi polynomials, *Memoirs Amer. Math. Soc.* **319** (1985) 1–55.
2. R. Askey and J. Wimp, Associated Laguerre and Hermite polynomials, *Proc. Roy. Soc. Edinburgh Sect. A* **96** (1984), 15–37.
3. W.N. Bailey, *Generalized Hypergeometric Series*, Cambridge Univ. Press, London, 1935.
4. J. Bustoz and M.E.H. Ismail, The associated ultraspherical polynomials and their q -analogues, *Canad. J. Math.* **34** (1982), 718–736.
5. G. Gasper and M. Rahman, *Basic Hypergeometric Series*, Cambridge Univ. Press, Cambridge, 1990.
6. W. Gautschi, Computational aspects of three-term recurrence relations, *SIAM Rev.* **9** (1967), 24–82.
7. D.P. Gupta and D.R. Masson, Exceptional q -Askey-Wilson polynomials and continued fractions, *Proc. A.M.S.* **112** (1991), 717–727.
8. D.P. Gupta, M.E.H. Ismail and D.R. Masson, Associated continuous Hahn polynomials, *Canad. J. of Math.* **43** (1991), 1263–1280.
9. D.P. Gupta, M.E.H. Ismail and D.R. Masson, Contiguous relations, Basic Hypergeometric functions and orthogonal polynomials II, Associated big q -Jacobi polynomials, *J. of Math. Analysis and Applications* **171** (1992), 477–497.
10. D.P. Gupta and D.R. Masson, Watson’s basic analogue of Ramanujan’s Entry 40 and its generalization, *SIAM J. Math. Anal.*, to appear.
11. M.E.H. Ismail, J. Letessier, and G. Valent, Linear birth and death models and associated Laguerre polynomials, *J. Approx. Theory* **56** (1988), 337–348.
12. M.E.H. Ismail, J. Letessier, and G. Valent, Quadratic birth and death processes and associated continuous dual Hahn polynomials, *SIAM J. Math. Anal.* **20** (1989), 727–737.
13. M.E.H. Ismail, J. Letessier, G. Valent and J. Wimp, Two families of associated Wilson polynomials, *Can. J. Math.* **42** (1990), 659–695.
14. M.E.H. Ismail and C.A. Libis, Contiguous relations, basic hypergeometric functions and orthogonal polynomials I, *J. Math. Anal. Appl.* **141** (1989), 349–372.
15. M.E.H. Ismail and D.R. Masson, Two families of orthogonal polynomials related to Jacobi polynomials, *Rocky Mountain J. Math.* **21** (1991), 359–375.
16. M.E.H. Ismail and M. Rahman, Associated Askey-Wilson polynomials, *Trans. Amer. Math. Soc.* **328** (1991), 201–239.

17. W.B. Jones and W.J. Thron, *Continued Fractions: Analytic Theory and Applications*, Addison-Wesley, Reading, Mass., 1980.
18. D.R. Masson, Associated Wilson polynomials, *Constructive Approximation* **7** (1991), 521–534.
19. ———, Wilson polynomials and some continued fractions of Ramanujan, *Rocky Mountain J. of Math.* **21** (1991), 489–499.
20. ———, The rotating harmonic oscillator eigenvalue problems, I. Continued fractions and analytic continuation, *J. Math. Phys.* **24** (1983), 2074–2088.
21. ———, Difference equations, continued fractions, Jacobi Matrices and orthogonal polynomials, In: *Non-linear numerical methods and Rational Approximation* (A. Cuyt, ed.) Dordrecht, Reidel, 1988, 239–257.
22. J.A. Wilson, Hypergeometric series, recurrence relations and some new orthogonal polynomials, Ph.D. diss., University of Wisconsin, Madison, 1978.
23. J. Wimp, Explicit formulas for the associated Jacobi polynomials and some applications, *Canad. J. Math.* **39** (1987), 983–1000.